

IMPACT OF ELECTRONIC HEALTH RECORD SYSTEMS ON QUALITY OF PRIMARY CARE WITHIN ESSENTIAL PUBLIC HEALTH FUNCTION FRAMEWORK

N. S. Yussupova^{1,2*}, B. S. Turdaliyeva¹, V. V. Koikov³, I.M. Son⁴, Zh. Y. Atalykova²

¹ Kazakh scientific center of dermatology and infectious diseases of the Ministry of Health of the Republic of Kazakhstan, Kazakhstan, Almaty

² NpJSC «Kazakh National Medical University named after S. D. Asfendiyarov», Kazakhstan, Almaty

³ NpJSC «Astana Medical University», Kazakhstan, Astana

⁴ Federal Research Institute for Health Organization and Informatics of the Ministry of Health of the Russian Federation, Russian, Moscow

*Corresponding author

Abstract

Quality healthcare is the primary objective of any healthcare system. An effective electronic health record system is a crucial tool for achieving this goal. Its objectives are to improve the quality of medical care by automating the work of doctors and medical personnel in all areas of activity.

The purpose of this study. To explore the impact of electronic health records systems in primary health care facilities on improving the quality of care to ensure a sustainable organizational structure as an essential public healthcare function.

Materials and methods. The search was conducted in the MEDLINE, EMBASE, CINAHL, and Cochrane Central Register of Controlled Trials databases. The search strategy was based on the PICO Framework. The studies selected for the meta-analysis were published between January 1, 2010, and May 1, 2023. Two authors independently reviewed article titles and abstracts for eligibility. Information from the search was deduplicated using EndNote X9 and imported into the Covidence Systematic Review for review. Statistical processing was performed in RStudio 2023.03.1 Build 446 (Posit Software, PBC).

Results. A literature search identified 640 publications; 11 of them were included in the review. Meta-analysis showed that the use of electronic health records (EHRs) helps to reduce the time to complete medical records by 33.4 % (95% CI = 0.8 % to 1.2 %; p < 0.007), promotes adherence to clinical recommendations (RR 1.30; 95 % CI = 1.04 to 1.79; p = 0.05) and reduces prescribing errors (RR 0.44; 95 % CI = 0.34 to 0.53; p < 0.001), which undoubtedly improves the quality of medical care.

Conclusion. This study validates the positive influence of EHRSs on enhancing care quality in PHC settings, primarily by streamlining documentation, reducing prescription errors, and aligning workflow with clinical guidelines. These findings can help healthcare and public health professionals make informed decisions about EHRS adoption.

Keywords: Electronic health record systems (EHRS), electronic health record (EHR), quality of care, sustainable organizational structure, meta-analysis, essential public health function (EPHF), primary health care (PHC).

Introduction

Quality healthcare is the primary objective of any healthcare system [1]. An effective electronic health record system (hereinafter – EHRS) is a key tool to achieve this goal. Its objectives are

to improve the quality of medical care by automating the work of doctors and medical personnel in all areas of activity, and to de-bureaucratize the work of medical staff [2]. This tool is recognized as one of the elements of a sustainable organizational

structure, which is also the fourth essential public health function of the health system, according to the World Health Organization document [3].

The World Health Assembly, in its resolution WHA69.1, identified the main public health functions as the most cost-effective, comprehensive, and sustainable ways to improve health and reduce disease burden [4].

The WHO Regional Office for Europe has developed and recommended ten essential public health functions, which serve as a detailed checklist of key public health activities and a resource for assessing public health services and capacities within a national approach. EPHFs are constantly evolving and should be updated regularly to reflect current assessments, new issues, and communication technologies, including social networks [5; 6].

The World Health Organization recognizes Primary Health Care (hereinafter – PHC) as the most important element of national health systems. WHO's policy of continuous improvement of PHC once again proves the essential contribution of PHC to individual and public health.

The following 5 functions are implemented at the PHC level:

1. Health promotion, including the impact on social determinants and reducing inequalities in health outcomes;
2. Prevention of diseases, including early detection of health problems;
3. Ensuring that the public health sector has sufficient qualified personnel;
4. Ensuring sustainable institutional structures and funding;
5. Advocacy, Communication, and Social Mobilization for Health.

Electronic health record systems (EHRSSs) ensure continuity of production and organizational processes by serving as direct elements of forming a stable organizational structure.

Increasingly, medical organizations are adopting electronic document processing to reduce the time required to complete necessary documentation [7]. At the same time, there is a lack of studies on the effectiveness of EHRSSs and their impact on the quality of care provided to ensure an organizational structure [8]. Contrary to the widespread belief that EHRSSs have an undeniable advantage, some studies show contradictory results, indicating

a decrease in the quality of medical care after the implementation of these systems [9; 10].

The purpose of this meta-analysis is to investigate the impact of EHRSSs on improving the quality of healthcare in order to ensure a sustainable organizational structure in PHC institutions, as an operational function of public health.

The study is of value to healthcare managers and public health professionals because the results can contribute to the optimization and improvement of the quality of medical care, including primary healthcare institutions.

The purpose of this study. To explore the impact of electronic health records systems in primary health care facilities on improving the quality of care to ensure a sustainable organizational structure as an essential public healthcare function.

Novelty of the work: For the first time, the impact of EHRSSs on improving the quality of medical care was examined to ensure a sustainable organizational structure in PHC institutions as an operational function of public health.

Materials and methods

Criteria for inclusion in the study: papers that explored the role of EHRS in improving quality in institutions with PHC units; papers examining the impact of EHRS on the quality of clinical care; research using electronic health record (hereinafter – EHR) data for decision making; works investigating the impact of EHRS on the organizational structure; studies published no earlier than January 01, 2010, and no later than May 01, 2023.

Criteria for exclusion from the study: works published not in the appropriate time interval; studies that are only available as abstracts or conference proceedings; studies whose conclusions are based on the study of patient feedback; studies that use performance data from third-party registries; studies in which EHRS resources were used to make decisions on the spot, without additional analysis of the information received.

Search strategy and study selection

The PICO Framework [11] guided our search strategy. EHRS refers to computerized patient data systems, including EHRs and clinical decision support systems (hereinafter – CDSS). Searches covered MEDLINE, EMBASE, CINAHL, and Cochrane databases, reflecting their scientific medical community use [12]. English language searches were used, including manual searches, citation searches, and

keywords: «Electronic health record», «Improving the quality of medical care», «Improvement», «Electronic decision support systems», «Optimization», and «Reduction of medical errors», «Electronic medical documentation», «Sustainable organizational structure», «Adverse drug reactions», «Completion of medical documentation».

Two authors independently screened article

titles and abstracts against exclusion criteria. Full texts were analyzed if exclusions were unclear. Disagreements were resolved by discussion. Full-text analysis was by one author, with inclusion confirmed by another. Our study adheres to PRISMA guidelines for systematic review and meta-analysis reporting [13]. The article selection process is illustrated in Figure 1.




Figure 1. Selection of studies for meta-analysis according to PRISMA guidelines [13]

Data extraction and quality assessment

The data collection template was developed in Microsoft Excel version 16.34. Using the Cochrane EPOC tool, the following data were extracted: authors' names, country, year, study design, conditions, duration, and results.

Statistical data processing

The search information was imported into the EndNote X9 link manager for deduplication and then imported into the Covidence Systematic Review software, a web-based review platform. Statistical processing was performed using RStudio 2023.03.1 Build 446 (Posit Software, PBC). A meta-analysis was conducted using a random effects model for each indicator. Heterogeneity was assessed using Cochran's Q-test.

Results

A literature search identified 640 studies.

After removing duplicates, 545 potentially relevant abstracts of articles were examined, and then 450 were excluded. A total of 95 full-text articles were selected for further review, of which 84 were excluded based on exclusion criteria. A total of 11 manuscripts were included in the review, which examined the relationship between the use of electronic medical records systems and the quality of care.

A summary of the main characteristics of the studies is described in Table 1. The design of the studies included in this meta-analysis was diverse, including review (n = 3), prospective cohort study (n = 1), systematic review (n = 1), descriptive qualitative study (n = 1), action research (n = 1), retrospective, observational study (n = 1), intervention evaluation study (n = 2), research before and after implementation (n=1).

Table 1. Main characteristics of the studies included in the meta-analysis

No.	Article	Study Design	Place of the study	Key parameter to include in meta-analysis	Indicators assessed during the meta-analysis	Average p-value for each indicator	Average confidence interval for each value
1	Robert S. Janet. 2019 [14]	Review	Multidisciplinary medical center with PHC departments and an emergency room (ER)	Application of EHRs	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	<0.007	0.8-1.2
2	Dara Koper [15]	Prospective cohort study	Multidisciplinary medical center with a PHC department	Application of EHRs	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	0.05	1.04-1.79
3	Sofia Säfholm [16]	Systematic review	For a systematic review, studies were taken that were conducted in multidisciplinary institutions with a PHC department	Application and implementation of EHRs	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	< 0.001	0.34-0.53
4	Kok Wai Kee [9]	Review	Medical centers (including PHC departments)	Consistent implementation of EHRs	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	< 0.001	0.43-0.90
5	Shamma Al Alawi [17]	Descriptive qualitative research	Health Center with the PHC Department	The use of paper carriers in comparison with EHRs	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	<0.001	61.0-79.4

6	Wetterneck TB et al., 2011 [18]	Research before and after implementation	City Medical Center at the University Hospital, including the Department of PHC	The effectiveness of the use of EHRS	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications	0.05 1.03-1.80
7	Shabbir SA et al., 2010 [19]	Action Research	Multidisciplinary medical center with a PHC department	Use of paper documentation and EHRS	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	0.05 1.04-1.33
8	Banerjee D., et al., 2017 [20]	Intervention Evaluation Study	Multidisciplinary medical center with a PHC department	Application of EHRS	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	0.03 1.04-1.75
9	Armor BL et al., 2016 [21]	Retrospective, observational study	Academic medical center with a PHC department	Prescribing medications EHRS	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	0.03 1.04-1.79
10	Chen J. et al., 2023 [22]	Intervention Evaluation Study	Oncology clinic with PHC department	Application of CDSS tools	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	0.05 1.03-1.67
11	Graeme N. et al., 2014 [7]	Review	For the review, studies were taken that were conducted in multidisciplinary institutions with a PHC department	EHR and EHRS	Time to complete medical records Adherence to clinical recommendations Mistakes in prescribing medications Adverse drug reactions	0.05 1.04-1.79

*Note: **EHRS** – Electronic Health Records System

ER – Emergency Room

EHR – Electronic Health Record

CDSS – Clinical Decision Support System

Source: elaborated by the authors.

Selected studies have examined the relationship between the use of EHRs and the reduction in time spent by healthcare professionals comple-

ting paperwork. The association between EHRs use and adherence, medication use, medical errors, and adverse effects was assessed in Table 2.

Table 2. Association between EHRs use and adherence, medication use, medical errors, and adverse effects

Index	Description	Value	95 % Confidence Interval	Significance level (<i>p</i>)
Time to complete documentation	%	33.4	0.8-1.2	<0.007
Adherence to clinical recommendations	Relative risk (RR)	1.30	1.04-1.79	0.05
Mistakes in prescribing medications	Relative risk (RR)	0.44	0.34-0.53	<0.001
Adverse drug reactions (ADRs)	Relative risk (RR)	0.59	0.43-0.90	0.045

Source: compiled by the authors

A meta-analysis found an association between the use of EHRS by medical professionals and a reduction in the time to complete documentation (Figure 2), with a mean difference of 33.4 % (95 % CI = 0.8 % to 1.2 %; *p* < 0.007). The use of EHR was also found to be associated with a higher rate of adherence to clinical guidelines (Figure 3a), a relative risk (RR) of 1.30 (95 % CI = 1.04 to 1.79; *p* = 0.05) and with a decrease in prescribing errors (Figure 3b), RR 0.44 (95 % CI = 0.34 to 0.53; *p* < 0.001), and with a reduction in adverse drug reac-

tions (Figure 3c), RR 0.59 (95 % CI = 0.43 to 0.90; *p* = 0.045). There was no association between EHR use and mortality (*p* = 0.936). High interstudy heterogeneity was found in terms of time spent completing documentation (Q-test *p* < 0, 001 and *I*² = 92.3 %), adherence to clinical guidelines (Q-test *p* < 0.001 and *I*² = 91.2 %), prescribing errors (Q-test *p* < 0.001 and *I*² = 96.1 %), and ADRs (Q-test *p* < 0.001 and *I*² = 79.4 %). There was also moderate heterogeneity in mortality (Q-test, *p* = 0.012; *I*² = 61.0 %).




Figure 2. Reduction of time costs when using EHRS

Source: compiled by the authors




Figure 3. Association of EHR use with (a) adherence to clinical guidelines, (b) reduction in prescribing errors, (c) reduction in adverse drug reactions

Source: compiled by the authors

Sensitivity analysis and publication bias

Sensitivity analysis showed that the overall effect remains stable even when any study is excluded from the analysis, and this does not lead to a significant improvement in the level of heterogeneity.

Discussion

The application of EHRS represents a significant step in improving healthcare effectiveness and sustainability [1; 2; 23].

By analyzing the meta-analysis results obtained in the context of ensuring a sustainable organizational structure, it is possible to highlight the significant role of EHRS in enhancing the quality of healthcare [16; 24].

Based on the results obtained in this study, we can affirm that the introduction and proper use of EHRS undoubtedly contribute to the improvement of the quality of medical care. These systems help to reduce the time (on average, 33.4%) that medical staff spend on completing paperwork, which in turn improves operational processes and allows staff to focus on more critical aspects of medical care. This represents a crucial balance between optimizing processes and maintaining a stable organizational structure.

An important aspect which is confirmed by this meta-analysis is the connection between the use of EHRS and the reduction of medication errors in prescribing medicines (RR 0.44; 95 % CI = 0.34 to 0.53; $p < .001$), as well as undesired drug

reactions (overall RR 0.59; 95 % CI = from 0.43 to 0.90; $p = 0.045$). It is known that effects like that can have a serious impact on patient health and the financial sustainability of medical institutions [25-27]. Reducing medical errors and adverse reactions from EHRS is important to patient safety and minimizing the cost of treating complications [28-30].

It is also important to consider that well-designed EHRS support adherence to clinical guidelines and protocols, which is important for maintaining the organizational structure [31-33]. This directly affects the operational function of public health, providing a more coherent and effective approach to treatment and care for patients [34; 35].

At the same time, in the clinical care setting, information is recorded into EHRs by various healthcare professionals, including physicians, nurses, medical assistants, and other clinical staff. The accuracy and completeness of the data entered depend heavily on the documentation practices of these health care providers, which can vary between institutions and countries.

Furthermore, we also face infrastructural problems, including accessibility, cost, and quality of internet in remote rural areas, as well as the availability of good computer technology.

Not all health managers and health workers are fully familiar with the digital tools available and the new opportunities they present. For many older doctors, the transition to EHRs is a challenge [36].

Nonetheless, we should note that despite its advantages, this study has limitations. Furthermore, increasing the heterogeneity between studies can affect the stability of results and requires further analysis. In addition, the technical aspects of different EHRSs are not always provided when evaluating a system, thereby limiting the full understanding and assessment of the effectiveness of these systems [37; 38].

Further studies devoted to technical aspects and implementation strategies will undoubtedly help improve and refine approaches to EHRS. Moreover, it will be possible to fully exploit their potential to maintain the effectiveness and stability of the health system.

Conclusion

A meta-analysis of studies examining the relationship between EHRS and quality improvement in healthcare, within the context of organizational sustainability, has provided valuable research results. A meta-analysis of sources, including 11 studies analyzed in detail, found statistically significant associations between the use of EHRS and improved quality of care. As a result, it became possible to formulate key recommendations for PHC medical institutions and the public health system.

A meta-analysis confirmed a significant association between the use of EHRS and a reduction in the time spent by medical professionals on completing documentation (mean reduction in time by 33.4 %, (95 % CI = 0.8 % to 1.2 %; p < 0.007). This is direct evidence of an increase in operational efficiency and the release of resources for more important medical tasks.

Most importantly, the use of EHRS was also associated with improved adherence to clinical guidelines (RR 1.30, 95 % CI 1.04-1.79; p = 0.05), highlighting the role of these systems in providing structured and coordinated medical care. Reduced prescribing errors (RR 0.44, 95 % CI = 0.34 to 0.53; p < 0.001) and adverse drug reactions (RR 0.59, 95 % CI = 0.43 to 0.90; p = 0.045) emphasize the importance of using EHRS to ensure patient safety and minimize adverse effects.

Despite the noted positive outcomes, it is crucial to acknowledge significant study heterogeneity, necessitating further scrutiny. These meta-analysis findings hold practical implications for healthcare providers, managers, and public health experts. The study confirms that implementing

electronic medical record systems enhances care quality, process efficiency, and organizational structure. More research on technical aspects and implementation strategies could deepen understanding of EHRS's practical role and potential in achieving a sustainable structure.

These results suggest the need for proactive implementation and optimization of EHRS in medical institutions to achieve enhanced care quality and a sustainable organizational structure. This approach, aligned with IT trends, will enhance treatment outcomes, resource utilization, patient safety, and organizational sustainability.

References

1. Pronovost P. J. High-Performing Health Care Delivery Systems: High Performance Toward What Purpose? // Jt Comm J Qual Patient Saf. – 2017. – Vol. 43(9). – P. 448-449. – DOI: 10.1016/j.jcqj.2017.06.001.
2. Jedwab R. M., Franco M., Owen D., Ingram A., Redley B., Dobroff N. Improving the Quality of Electronic Medical Record Documentation: Development of a Compliance and Quality Program // Apple Clinic Inform. – 2022. – Vol. 13(4). – P. 836-844. – DOI: 10.1055/s-0042-1756369.
3. World Health Organization. Essential public health function Suggested citation. ns, health systems and health security: developing conceptual clarity and a WHO roadmap for action. Geneva: World Health Organization. – 2018. – 67 p.
4. World Health Organization. SIXTY-NINTH WORLD HEALTH ASSEMBLY. WHA69.1. Resolution adopted by the World Health Assembly. Geneva: World Health Organization. – 2016. – 450 p.
5. Слабкий Г. А., Пархоменко Г. Я. Міжнародні підходи до розвитку громадського здоров'я» // Україна. Здоров'я Нації. – 2012. – № 1(21). – C. 7.
6. Strengthening Public Health Services in the European Region – Brief Overview of Source Documents for the European Action Plan [Electronic source] // World health organization [website]. – URL: http://www.euro.who.int/_data/assets/pdf_file/0007/172681/RC62-id05-final-Rus.pdf?ua=1 (Accessed: 06.02.2025).
7. Graeme N. Forrest and others, Use of Electronic Health Records and Clinical Decision Support Systems for Antimicrobial Stewardship // Clinical In-

- fectious Diseases. – Vol. 59. – 2014. – P. 122-133. – DOI: <https://doi.org/10.1093/cid/ciu565>.
8. Joyst K. E., Bhatt D. L., Schwamm L. H., Xian Y., Heidenreich P. A., Fonarow G. C., Smith E. E., Neely M. L., Grau-Sepulveda M. V., Hernandez A. F. Lack of impact of electronic health records on quality of care and outcomes for ischemic stroke // *J Am Coll Cardiol.* – 2015. – Vol. 65(18). – P. 1964-1972. – DOI: 10.1016/j.jacc.2015.02.059.
9. Kee K. W., Char C. W., Yip A. Y. A review on interventions to reduce medication discrepancies or errors in primary or ambulatory care setting during care transition from hospital to primary care // *J Family Med Prim Care.* – 2018. – Vol. 7. – P. 501-506.
10. Tsai C. H., Eghdam A., Davoody N., Wright G., Flowerday S., Koch S. Effects of Electronic Health Record Implementation and Barriers to Adoption and Use: A Scoping Review and Qualitative Analysis of the Content. *Life (Basel).* – 2020. – Vol. 10(12). – P. 327. – DOI: 10.3390/life10120327.
11. Agoritsas T., Merglen A., Courvoisier D.S., Combescure C., Garin N., Perrier A., Perneger T. V. Sensitivity and predictive value of 15 PubMed search strategies to answer clinical questions rated against full systematic reviews // *J Med Internet Res.* – 2012. – Vol. 14(3). – P. 85. – DOI: 10.2196/jmir.2021.
12. Kwon Y., Lemieux M., McTavish J., Wathen N. Identifying and removing duplicate records from systematic review searches // *J Med Libr Assoc.* – 2015. – Vol. 103(4). – P. 184-188. – DOI: 10.3163/1536-5050.103.4.004.
13. Page M. J., McKenzie J. E., Bossuyt P. M., Boutron I., Hoffmann T. C., Mulrow C. D., Shamseer L., Tetzlaff J. M., Akl E. A., Brennan S. E., Chou R., Glanville J., Grimshaw J. M., Hróbjartsson A., Lalu M. M., Li T., Loder E. W., Mayo-Wilson E., McDonald S., McGuinness L. A., Stewart L. A., Thomas J., Tricco A. C., Welch V. A., Whiting P., Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews // *BMJ.* – 2021. – Vol. 372. – P. 71. – DOI: 10.1136/bmj.n71.
14. Janett R. S., Yeracaris P. P. Electronic Medical Records in the American Health System: challenges and lessons learned // *Cien Saude Colet.* – 2020. – Vol. 25(4). – P. 1293-1304. – DOI: 10.1590/1413-81232020254.28922019.
15. Koper D., Kamenski G., Flamm M., Böhmdorfer B. Frequency of medication errors in primary care patients with polypharmacy // *Family Practice.* – 2013. – Vol. 30(3). – P. 313-319. – DOI: <https://doi.org/10.1093/fampra/cms070>.
16. Säfholm S., Bondesson Å. & Modig, S. Medication errors in primary health care records; a cross-sectional study in Southern Sweden // *BMC Fam Pract.* – 2019. – Vol. 20. – P. 110. – DOI: <https://doi.org/10.1186/s12875-019-1001-0>.
17. Al Alawi S., Al Dhaheri A., Al Baloushi D. et al. Physician user satisfaction with an electronic medical records system in primary healthcare centers in Al Ain: a qualitative study // *BMJ Open.* – 2014. – Vol. 4(11). – Article No. 005569. – DOI: 10.1136/bmjopen-2014-005569.
18. Wetterneck T. B., Walker J. M., Blosky M. A., Cartmill R. S., Hoonakker P., Johnson M. A., Norfolk E., Carayon P. Factors contributing to an increase in duplicate medication order errors after CPOE implementation // *J Am Med Inform Assoc.* – 2011. – Vol. 18(6). – P. 774-782. – DOI: 10.1136/amiajnl-2011-000255.
19. Shabbir S. A., Ahmed L. A., Sudhir R. R., Scholl J., Li Y. C., Liou D. M. Comparison of documentation time between an electronic and a paper-based record system by optometrists at an eye hospital in south India: a time-motion study // *Comput Methods Programs Biomed.* – 2010. – Vol. 100(3). – P. 283-288. – DOI: 10.1016/j.cmpb.2010.04.003.
20. Banerjee D., Thompson C., Kell C., Shetty R., Vetteth Y., Grossman H., DiBiase A., Fowler M. An informatics-based approach to reducing heart failure all-cause readmissions: the Stanford heart failure dashboard // *J Am Med Inform Assoc.* – 2017. – Vol. 24(3). – P. 550-555. – DOI: 10.1093/jamia/ocw150.
21. Armor B. L., Wight A. J., Carter S.M. Evaluation of Adverse Drug Events and Medication Discrepancies in Transitions of Care Between Hospital Discharge and Primary Care Follow-Up // *J Pharm Pract.* – 2016. – Vol. 29(2). – P. 132-137. – DOI: 10.1177/0897190014549836.
22. Chen J., Cutrona S. L., Dharod A., Bunch S. C., Foley K. L., Ostasiewski B., Hale E. R., Bridges A., Moses A., Donny E. C., Sutfin E. L., Houston T. K. iDAPT Implementation Science Center for Cancer Control. Monitoring the Implementation of Tobacco Cessation Support Tools: Using Novel Electronic Health Record Activity Metrics // *JMIR MedInform.* – 2023. – Vol. 11. – Article No. 43097. – DOI: 10.2196/43097.

23. Pohlmann S., Kunz A., Ose D., Winkler E. C., Brandner A., Poss-Doering R., Szecsenyi J., Wensing M. Digitalizing Health Services by Implementing a Personal Electronic Health Record in Germany: Qualitative Analysis of Fundamental Prerequisites from the Perspective of Selected Experts // *J Med Internet Res.* – 2020. – Vol. 22(1). – Article No. 15102. – DOI: 10.2196/15102.
24. Robinson K. E., Kersey J. A. Novel electronic health record (EHR) education intervention in large healthcare organization improves quality, efficiency, time, and impact on burnout // *Medicine (Baltimore)*. – 2018. – Vol. 97(38). – Article No. 12319. – DOI: 10.1097/MD.00000000000012319.
25. Shafi S., Collinsworth A. W., Copeland L. A., Ogola G. O., Qiu T., Kouznetsova M., Liao I. C., Mears N., Pham A.T., Wan G. J., Masica A. L. Association of Opioid-Related Adverse Drug Events with Clinical and Cost Outcomes Among Surgical Patients in a Large Integrated Health Care Delivery System // *JAMA Surg.* – 2018. – Vol. 153(8). – P. 757-763. – DOI: 10.1001/jamasurg.2018.1039.
26. Liao P. J., Mao C. T., Chen T. L., Deng S. T., Hsu K. H. Factors associated with adverse drug reaction occurrence and prognosis, and their economic impacts in older inpatients in Taiwan: a nested case-control study // *BMJ Open*. – 2019. – Vol. 9(5). – Article No. 026771. – DOI: 10.1136/bmjjopen-2018-026771.
27. Yagi M., Shindo Y., Mutoh Y., Sano M., Sakakibara T., Kobayashi H., Matsuura A., Emoto R., Matsui S., Nakagawa T., Ogawa K. Factors associated with adverse drug reactions or death in very elderly hospitalized patients with pulmonary tuberculosis // *Scientific Reports*. – 2023. – Vol. 13(1). – Article No. 6826. – DOI: 10.1038/s41598-023-33967-6.
28. de Hoon SEM, Hek K., van Dijk L., Verheij R.A. Adverse events recording in electronic health record systems in primary care // *BMC Med Inform Decis Mak.* – 2017. – Vol. 17(1). – P. 163. – DOI: 10.1186/s12911-017-0565-7.
29. Zarrinpar A., David Cheng T. Y., Huo Z. What Can We Learn About Drug Safety and Other Effects in the Era of Electronic Health Records and Big Data That We Would Not Be Able to Learn from Classic Epidemiology? // *J Surg Res.* – 2020. – Vol. 246. – P. 599-604. – DOI: 10.1016/j.jss.2019.09.053.
30. Hui C., Vaillancourt R., Bair L., Wong E., King J.W. Accuracy of Adverse Drug Reaction Documentation upon Implementation of an Ambulatory Electronic Health Record System // *Drugs Real World Outcomes*. – 2016. – Vol. 3(2). – P. 231-238. – DOI: 10.1007/s40801-016-0071-8.
31. Kern L. M., Barron Y., Dhopeshwarkar R. V., Edwards A., Kaushal R. HITEC Investigators. Electronic health records and ambulatory quality of care // *JG Intern Med.* – 2013. – Vol. 28(4). – P. 496-503. – DOI: 10.1007/s11606-012-2237-8.
32. Personal Electronic Health Records: A Review of Clinical Effectiveness, Cost-Effectiveness, and Guidelines [Electronic source] // Ottawa (ON): Canadian Agency for Drugs and Technologies in Health CADTH Rapid Response Reports [website]. – 2016. – URL: <https://www.ncbi.nlm.nih.gov/books/NBK355652/> (Accessed: 10.02.2025).
33. 21st century health challenges: can the essential public health functions make a difference? [Electronic source]. – Geneva: World Health Organization. – 2021. – 90 p. – URL: <https://www.who.int/publications/item/9789240038927>. (Accessed: 12.02.2025)
34. van Fenema E. M. Meten van naleving van richtlijnen en behandelkwaliteit met data van routine outcome monitoring // *Tijdschrift voor Psychiatrie*. – 2017. – Vol. 59(3). – P. 159-165.
35. Cowppli-Bony A., Tretarre B., Marrer E., Defossez G., Daubisse-Marliac L., Coureau G., Minicozzi P., Woronoff A.S., Delafosse P., Molinié F. FRANCIM network. Compliance with clinical guidelines for breast cancer management: A population-based study of quality-of-care indicators in France // *PLOS One*. – 2019. – Vol. 14(10). – Article No. 0224275. – DOI: 10.1371/journal.pone.0224275.
36. Nijor S., Rallis G., Lad N., Gokcen E. Patient safety issues from information overload in electronic medical records // *J Patient Saf.* – 2022. – Vol. 18(6). – P. 999-1003.
37. Fennelly O., Cunningham C., Grogan L., Cronin H., O’Shea C., Roche M., Lawlor F., O’Hare N. Successfully implementing a national electronic health record: a rapid umbrella review // *Int J Med Inform.* – 2020. – Vol. 144. – Article No. 104281. – DOI: 10.1016/j.ijmedinf.2020.104281.
38. Holmes J. H., Beinlich J., Boland M. R., Bowles K. H., Chen Y., Cook T. S., Demiris G., Draughelis M., Fluharty L., Gabriel P.E., Grundmeier R., Hanson C.W., Herman D. S., Himes B. E., Hubbard R. A., Kahn C. E. Jr, Kim D., Koppel R., Long Q., Mirkovic N., Morris J. S., Mowery D. L., Ritchie

M. D., Urbanowicz R., Moore J. H. Why Is the Electronic Health Record So Challenging for Research and Clinical Care? // Methods Inf Med. – 2021. – Vol. 60(1-02). – P. 32-48. – DOI: 10.1055/s-0041-1731784.

References

1. Pronovost, P. J. (2017). High-performing health care delivery systems: High performance toward what purpose? *Joint Commission Journal on Quality and Patient Safety*, 43(9), 448-449, DOI: <https://doi.org/10.1016/j.jcjq.2017.06.001>
2. Jedwab, R. M., Franco, M., Owen, D., Ingram, A., Redley, B., & Dobroff, N. (2022). Improving the quality of electronic medical record documentation: Development of a compliance and quality program. *Applied Clinical Informatics*, 13(4), 836–844. DOI: <https://doi.org/10.1055/s-0042-1756369>.
3. World Health Organization (2018). Essential public health functions, health systems and health security: Developing conceptual clarity and a WHO roadmap for action. World Health Organization, 67 p.
4. World Health Organization (2016). Sixty-ninth World Health Assembly: WHA69.1. Resolution adopted by the World Health Assembly. World Health Organization, 450 p.
5. Slabkij, G. A., & Parhomenko, G. Ja. (2012). Mizhnarodni pidhodi do rozvitku gromads'kogo zdrorov'ja. Ukraine. *Health of the nation*, 21(1), 7.
6. Strengthening public health services in the European Region – Brief overview of source documents for the European action plan. World Health Organization [website]. Retrieved February 6, 2025, from http://www.euro.who.int/_data/assets/pdf_file/0007/172681/RC62-id05-final-Rus.pdf?ua=1.
7. Forrest, G. N., et al. (2014). Use of electronic health records and clinical decision support systems for antimicrobial stewardship. *Clinical Infectious Diseases*, 59, 122-133. DOI:<https://doi.org/10.1093/cid/ciu565>
8. Joynt, K. E., Bhatt, D. L., Schwamm, L. H., Xian, Y., Heidenreich, P. A., Fonarow, G. C., Smith, E. E., Neely, M. L., Grau-Sepulveda, M. V., & Hernandez, A. F. (2015). Lack of impact of electronic health records on quality of care and outcomes for ischemic stroke. *Journal of the American College of Cardiology*, 65(18), 1964-1972. DOI: <https://doi.org/10.1016/j.jacc.2015.02.059>.
9. Kee, K. W., Char, C. W., & Yip, A. Y. (2018). A review on interventions to reduce medication discrepancies or errors in primary or ambulatory care setting during care transition from hospital to primary care. *Journal of Family Medicine and Primary Care*, 7, 501-506.
10. Tsai, C. H., Eghdam, A., Davoodi, N., Wright, G., Flowerday, S., & Koch, S. (2020). Effects of electronic health record implementation and barriers to adoption and use: A scoping review and qualitative analysis of the content. *Life (Basel)*, 10(12), 327. DOI: <https://doi.org/10.3390/life10120327>.
11. Agoritsas, T., Merglen, A., Courvoisier, D. S., Combescure, C., Garin, N., Perrier, A., & Perneger, T. V. (2012). Sensitivity and predictive value of 15 PubMed search strategies to answer clinical questions rated against full systematic reviews. *Journal of Medical Internet Research*, 14(3), 85. DOI: <https://doi.org/10.2196/jmir.2021>.
12. Kwon, Y., Lemieux, M., McTavish, J., & Wathen, N. (2015). Identifying and removing duplicate records from systematic review searches. *Journal of the Medical Library Association*, 103(4), 184-188. DOI: <https://doi.org/10.3163/1536-5050.103.4.004>.
13. Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., et al. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 372, 71. DOI: <https://doi.org/10.1136/bmj.n71>
14. Janett, R. S., & Yeracaris, P. P. (2020). Electronic medical records in the American health system: Challenges and lessons learned. *Ciência & Saúde Coletiva*, 25(4), 1293-1304. DOI: <https://doi.org/10.1590/1413-81232020254.28922019>.
15. Koper, D., Kamenski, G., Flamm, M., & Böhmdorfer, B. (2013). Frequency of medication errors in primary care patients with polypharmacy. *Family Practice*, 30(3), 313-319. DOI: <https://doi.org/10.1093/fampra/cms070>.
16. Säfholm, S., Bondesson, Å., & Modig, S. (2019). Medication errors in primary health care records: A cross-sectional study in Southern Sweden. *BMC Family Practice*, 20, 110. DOI: <https://doi.org/10.1186/s12875-019-1001-0>.
17. Al Alawi, S., Al Dhaheri, A., Al Baloushi, D., et al. (2014). Physician user satisfaction with an electronic medical records system in primary healthcare centers in Al Ain: A qualitative study. *BMJ Open*, 4(11), e005569. DOI: <https://doi.org/10.1136/bmjjopen-2014-005569>.

18. Wetterneck, T. B., Walker, J. M., Bloksy, M. A., Cartmill, R. S., Hoonakker, P., Johnson, M. A., Norfolk, E., & Carayon, P. (2011). Factors contributing to an increase in duplicate medication order errors after CPOE implementation. *Journal of the American Medical Informatics Association*, 18(6), 774-782. DOI: <https://doi.org/10.1136/amia-jnl-2011-000255>.
19. Shabbir, S. A., Ahmed, L. A., Sudhir, R. R., Scholl, J., Li, Y. C., & Liou, D. M. (2010). Comparison of documentation time between an electronic and a paper-based record system by optometrists at an eye hospital in South India: A time-motion study. *Computer Methods and Programs in Biomedicine*, 100(3), 283-288. DOI: <https://doi.org/10.1016/j.cmpb.2010.04.003>.
20. Banerjee, D., Thompson, C., Kell, C., Shetty, R., Vetteth, Y., Grossman, H., DiBiase, A., & Fowler, M. (2017). An informatics-based approach to reducing heart failure all-cause readmissions: The Stanford heart failure dashboard. *Journal of the American Medical Informatics Association*, 24(3), 550-555. DOI: <https://doi.org/10.1093/jamia/ocw150>.
21. Armor, B. L., Wight, A. J., & Carter, S. M. (2016). Evaluation of adverse drug events and medication discrepancies in transitions of care between hospital discharge and primary care follow-up. *Journal of Pharmacy Practice*, 29(2), 132-137. DOI: <https://doi.org/10.1177/0897190014549836>.
22. Chen, J., Cutrona, S. L., Dharod, A., Bunch, S. C., Foley, K. L., Ostasiewski, B., Hale, E. R., Bridges, A., Moses, A., Donny, E. C., Sutfin, E. L., & Houston, T. K. (2023). Monitoring the implementation of tobacco cessation support tools: Using novel electronic health record activity metrics. *JMIR Medical Informatics*, 11, e43097. DOI: <https://doi.org/10.2196/43097>.
23. Pohlmann, S., Kunz, A., Ose, D., Winkler, E. C., Brandner, A., Poss-Doering, R., Szecsenyi, J., & Wensing, M. (2020). Digitalizing health services by implementing a personal electronic health record in Germany: Qualitative analysis of fundamental prerequisites from the perspective of selected experts. *Journal of Medical Internet Research*, 22(1), e15102. DOI: <https://doi.org/10.2196/15102>
24. Robinson, K. E., & Kersey, J. A. (2018). Novel electronic health record (EHR) education intervention in large healthcare organization improves quality, efficiency, time, and impact on burnout. *Medicine*, 97(38), e12319. DOI: <https://doi.org/10.1097/MD.00000000000012319>.
25. Shafi, S., Collinsworth, A. W., Copeland, L. A., Ogola, G. O., Qiu, T., Kouznetsova, M., Liao, I. C., Mears, N., Pham, A. T., Wan, G. J., & Masica, A. L. (2018). Association of opioid-related adverse drug events with clinical and cost outcomes among surgical patients in a large integrated health care delivery system. *JAMA Surgery*, 153(8), 757-763. DOI: <https://doi.org/10.1001/jamasurg.2018.1039>
26. Liao, P. J., Mao, C. T., Chen, T. L., Deng, S. T., & Hsu, K. H. (2019). Factors associated with adverse drug reaction occurrence and prognosis, and their economic impacts in older inpatients in Taiwan: A nested case-control study. *BMJ Open*, 9(5), e026771. DOI: <https://doi.org/10.1136/bmjopen-2018-026771>.
27. Yagi, M., Shindo, Y., Mutoh, Y., Sano, M., Sakakibara, T., Kobayashi, H., Matsuura, A., Emoto, R., Matsui, S., Nakagawa, T., & Ogawa, K. (2023). Factors associated with adverse drug reactions or death in very elderly hospitalized patients with pulmonary tuberculosis. *Scientific Reports*, 13(1), 6826. DOI: <https://doi.org/10.1038/s41598-023-33967-6>.
28. de Hoon, S. E. M., Hek, K., van Dijk, L., & Verheij, R. A. (2017). Adverse events recording in electronic health record systems in primary care. *BMC Medical Informatics and Decision Making*, 17(1), 163. DOI: <https://doi.org/10.1186/s12911-017-0565-7>.
29. Zarrinpar, A., Cheng, T. Y. D., & Huo, Z. (2020). What can we learn about drug safety and other effects in the era of electronic health records and big data that we would not be able to learn from classic epidemiology? *Journal of Surgical Research*, 246, 599-604. DOI: <https://doi.org/10.1016/j.jss.2019.09.053>.
30. Hui, C., Vaillancourt, R., Bair, L., Wong, E., & King, J. W. (2016). Accuracy of adverse drug reaction documentation upon implementation of an ambulatory electronic health record system. *Drugs – Real World Outcomes*, 3(2), 231-238. DOI: <https://doi.org/10.1007/s40801-016-0071-8>.
31. Kern, L. M., Barron, Y., Dhopeshwarkar, R. V., Edwards, A., & Kaushal, R. (2013). Electronic health records and ambulatory quality of care. *Journal of General Internal Medicine*, 28(4), 496-503. DOI: <https://doi.org/10.1007/s11606-012-2237-8>.
32. Personal Electronic Health Records: A Review of Clinical Effectiveness, Cost-Effectiveness, and



- Guidelines (2016). Ottawa (ON): Canadian Agency for Drugs and Technologies in Health CADTH Rapid Response Reports [website]. Retrieved February 10, 2025, from <https://www.ncbi.nlm.nih.gov/books/NBK355652/>.
33. 21st century health challenges: Can the essential public health functions make a difference? (2021). World Health Organization [website]. Retrieved February 12, 2025, from <https://www.who.int/publications/i/item/9789240038927>.
34. van Fenema, E. M. (2017). Meten van naleving van richtlijnen en behandelkwaliteit met data van routine outcome monitoring [Assessment of guideline adherence and quality of care with routine outcome monitoring data]. *Journal of Psychiatry*, 59(3), 159-165.
35. Cowppli-Bony, A., Tretarre, B., Marrer, E., Defossez, G., Daubisse-Marliac, L., Coureau, G., Minicozzi, P., Woronoff, A. S., Delafosse, P., & Molinié, F. (2019). Compliance with clinical guidelines for breast cancer management: A population-based study of quality-of-care indicators in France. *PLOS ONE*, 14(10), e0224275. DOI: <https://doi.org/10.1371/journal.pone.0224275>.
36. Nijor, S., Rallis, G., Lad, N., & Gokcen, E. (2022). Patient safety issues from information overload in electronic medical records. *Journal of Patient Safety*, 18(6), 999-1003.
37. Fennelly, O., Cunningham, C., Grogan, L., Cronin, H., O'Shea, C., Roche, M., Lawlor, F., & O'Hare, N. (2020). Successfully implementing a national electronic health record: A rapid umbrella review. *International Journal of Medical Informatics*, 144, 104281. DOI <https://doi.org/10.1016/j.ijmedinf.2020.104281>
38. Holmes, J. H., Beinlich, J., Boland, M. R., Bowles, K. H., Chen, Y., Cook, T. S., Demiris, G., Draugelis, M., Fluharty, L., Gabriel, P. E., Grundmeier, R., Hanson, C. W., Herman, D. S., Himes, B. E., Hubbard, R. A., Kahn, C. E., Kim, D., Koppel, R., Long, Q., Mirkovic, N., Morris, J. S., Mowery, D. L., Ritchie, M. D., Urbanowicz, R., & Moore, J. H. (2021). Why is the electronic health record so challenging for research and clinical care? *Methods of Information in Medicine*, 60(1-2), 32-48. DOI: <https://doi.org/10.1055/s-0041-1731784..>

АЛҒАШҚЫ МЕДИЦИНАЛЫҚ-САНИТАРЛЫҚ КӨМЕК ДЕНГЕЙІНДЕ ҚОҒАМДЫҚ ДЕНСАУЛЫҚ САҚТАУДЫҢ ОПЕРАТИВТІ ФУНКЦИЯСЫН ҚАМТАМАСЫЗ ЕТУ ШЕҢБЕРІНДЕ МЕДИЦИНАЛЫҚ ҚҰЖАТТАМАНЫҢ ЭЛЕКТРОНДЫҚ ЖҮЙЕЛЕРИНІҢ МЕДИЦИНАЛЫҚ КӨМЕК САПАСЫНА ӘСЕРІ

Н. С. Юсупова^{1,2*}, Б. С. Турдалиева¹, В. В. Койков³, И. М. Соң⁴, Ж. Е. Аталькова²

¹ ҚР ДСМ Қазақ Дерматология және инфекциялық аурулар ғылыми орталығы»,
Қазақстан, Алматы

² «С. Д. Асфендияров атындағы Қазақ Ұлттық медицина университеті» КеАҚ,
Қазақстан, Алматы

³ «Астана медицина университеті» КеАҚ, Қазақстан, Астана

⁴ Ресей Федерациясы Денсаулық сақтау министрлігінің «Орталық ғылыми-зерттеу денсаулық
сақтауды үйымдастыру және ақпараттандыру институты»
Федералды мемлекеттік бюджеттік мекемесі, Ресей, Мәскеу

*Корреспондент автор

Андатпа

Сапалы медициналық көмек көрсету-кез келген денсаулық сақтау жүйесінің негізгі мақсаты. Мақсатқа жетудің негізгі құралдарының бірі-медициналық құжаттаманың тиімді жүйесі. Оның міндеттері-қызметтің барлық бағыттары бойынша дәрігерлер мен медициналық персоналдың жұмысын автоматтандыру есебінен медициналық көмектің сапасын арттыру.

Мақсаты. Қоғамдық денсаулық сақтаудың жедел функциясы ретінде тұрақты ұйымдық құрылымды қамтамасыз ету мақсатында алғашқы медициналық-санитарлық көмек мекемелеріндегі медициналық құжаттаманың электрондық жүйелерінің медициналық көмектің сапасын арттыруға әсерін зерттеу.

Материалдар мен әдістер. Іздеу MEDLINE, EMBASE, CINAHL дерекқорларында және Cochrane бақыланатын сынақтардың орталық тізлімінде жүргізілді. Іздеу стратегиясы PICO Framework-қа негізделген. Мета-талдау үшін таңдалған зерттеулер 01.01.2010 және 01.05.2023 жылдар аралығында жарияланды. Екі автор мақалалардың тақырыптары мен критерийлерге сәйкестігін тәуелсіз зерттеді. Іздеуден алынған ақпарат EndNote X9 көмегімен көшірілді, текстеру үшін Covid Systematic Review-ке импортталды. Статистикалық өңдеу R Studio 2023.03.1 Build 446 (Post Software, PC) бағдарламасында орындалды. Мета-талдау әр көрсеткіш үшін кездескөн эффект моделін қолдана отырып жүргізілді, гетерогенділік Кохреннің Q критерийі арқылы бағаланды.

Нәтижелер. Әдебиеттерді іздеу 640 зерттеуді анықтады, шолуға 11 қолжазба енгізілді. Мета-талдау көрсеткендей, электронды медициналық карталарды (ЕМС) пайдалану медициналық құжаттарды толтыру уақытын 33,4 %-ға (95 % ci = 0,8 %-дан 1,2 %-ға дейін; p < 0,007) қысқартуға көмектеседі, клиникалық ұсыныстардың сақталуына ықпал етеді (o 1,30; 95 % ci = 1,04-тен 1,79-ға дейін; p = 0,05) және дәрі-дәрмектерді тағайындау кезінде қателерді азайтуға мүмкіндік береді (o 0,44; 95 % ci = 0,34 0,53 дейін; p < 0,001), бұл сөзсіз медициналық көмектің сапасын арттырады.

Қорытынды. Бұл зерттеу медициналық құжаттаманың электрондық жүйелерінің алғашқы медициналық-санитариялық көмек мекемелеріндегі медициналық көмектің сапасын арттыруға, негізінен құжаттаманы толтыру уақытын оңтайландыру, персоналдың қателер санын азайту және клиникалық ұсынымдарға сәйкес жұмысты ұйымдастыру есебінен оң әсерін растайды. Зерттеу нәтижелері медициналық құжаттаманың электрондық жүйелерін енгізу туралы негізделген шешімдер қабылдау кезінде денсаулық сақтауды ұйымдастыру мамандары, қоғамдық денсаулық сақтау мамандары үшін пайдалы болады.

Түйін сөздер: медициналық құжаттаманың электрондық жүйелері (МКЭЖ), электрондық медициналық карта (ЭМК), медициналық көмектің сапасы, тұрақты Ұйымдық құрылым, мета-талдау, оперативті функция.

ВЛИЯНИЕ ЭЛЕКТРОННЫХ СИСТЕМ МЕДИЦИНСКОЙ ДОКУМЕНТАЦИИ НА КАЧЕСТВО МЕДИЦИНСКОЙ ПОМОЩИ В РАМКАХ ОБЕСПЕЧЕНИЯ ОПЕРАТИВНОЙ ФУНКЦИИ ОБЩЕСТВЕННОГО ЗДРАВООХРАНЕНИЯ НА УРОВНЕ ПМСП

Н. С. Юсупова^{1,2*}, Б. С. Турдалиева¹, В. В. Койков³, И. М. Сон⁴, Ж. Е. Аталькова²

¹ РГП на ПХВ «Казахский научный центр дерматологии и инфекционных заболеваний»

Министерства здравоохранения Республики Казахстан, Казахстан, Алматы

² НАО «Казахский Национальный Медицинский Университет имени С. Д. Асфендиярова»,

Казахстан, Алматы

³ НАО «Медицинский университет Астана», Казахстан, Астана

⁴ Федеральное государственное бюджетное учреждение «Центральный научно-исследовательский институт организации и информатизации здравоохранения»

Министерства здравоохранения Российской Федерации, Россия, Москва

*Корреспондирующий автор

Аннотация

Оказание качественной медицинской помощи – основополагающая цель любой системы здравоохранения. Одним из ключевых инструментов для достижения цели является эффективная система медицинской документации. Ее задачи – повышение качества медицинской помощи за счет автоматизации работы врачей и медицинского персонала по всем направлениям деятельности.

Цель. Изучить влияние электронных систем медицинской документации в учреждениях первичной медико-санитарной помощи на повышение качества медицинской помощи в целях обеспечения устойчивой организационной структуры, как оперативной функции общественного здравоохранения.

Материалы и методы. Поиск осуществлялся в базах данных MEDLINE, EMBASE, CINAHL и в Kokranovском центральном регистре контролируемых испытаний. Стратегия поиска базирова-



лась на PICO Framework. Исследования, отобранные для мета-анализа, были опубликованы в период с 01.01.2010 г. по 01.05.2023 г. Два автора независимо друг от друга изучали заголовки и резюме статей на соответствие критериям. Информация из поиска была дедублицирована с помощью EndNote X9, импортирована в Covidence Systematic Review для проверки. Статистическую обработку выполняли в RStudio 2023.03.1 Build 446 (Posit Software, РВС). Мета-анализ проводили с использованием модели случайных эффектов для каждого показателя, гетерогенность оценивали через Q-критерий Кохрена.

Результаты. Поиск литературы выявил 640 публикаций, 11 статей были включены в обзор. Мета-анализ показал, что использование электронных медицинских карт (ЭМК) помогает сократить время на заполнение медицинской документации на 33,4 % (95 % ДИ = 0,8 % до 1,2 %; $p < 0,007$), способствует соблюдению клинических рекомендаций (ОР 1,30; 95 % ДИ = 1,04 до 1,79; $p = 0,05$) и позволяет снизить ошибки при назначении препаратов (ОР 0,44; 95 % ДИ = 0,34 до 0,53; $p < 0,001$), что несомненно повышает качество медицинской помощи.

Заключение. Данное исследование подтверждает положительное влияние электронных систем медицинской документации на повышение качества медицинской помощи в учреждениях первичной медико-санитарной помощи, главным образом за счет оптимизации времени заполнения документации, снижения количества ошибок персонала и организации работы в соответствии с клиническими рекомендациями. Результаты исследования будут полезными для специалистов по организации здравоохранения, специалистов общественного здравоохранения при принятии обоснованных решений о внедрении электронных систем медицинской документации.

Ключевые слова: Электронные системы медицинской документации (ЭСМД), электронная медицинская карта (ЭМК), качество медицинской помощи, устойчивая организационная структура, мета-анализ, оперативная функция.

АВТОРЛАР ТУРАЛЫ

Юсупова Наргиза Сайдахметовна – медицина ғылымдарының магистрі, халықаралық ынтымақтастық жөніндегі менеджер, КР ДСМ «Қазақ Dermatология және инфекциялық аурулар ғылыми орталығы»; «Қоғамдық денсаулық сақтау» мамандығы бойынша докторантурасы бағдарламасының түлегі, «С. Д. Асфендияров атындағы Қазақ Ұлттық медицина университеті» КеАҚ; e-mail: ns.yussupova@gmail.com; ORCID:0000-0003-1857-2800.

Турдалиева Ботагоз Сайтовна – медицина ғылымдарының докторы, профессор, ғылыми қызмет және стратегиялық даму бойынша директордың орынбасары, КР ДСМ «Қазақ Dermatология және инфекциялық аурулар ғылыми орталығы»; e-mail: botturd@gmail.com; ORCID:0000-0001-9884-0777.

Койков Виталий Викторович – медицина ғылымдарының докторы, проректор, «Астана медицина университеті» КеАҚ; e-mail: koykov@inbox.ru; ORCID:0000-0001-7668-5614.

Сон Ирина Михайловна – медицина ғылымдарының докторы, профессор, Ресей Федерациясы Денсаулық сақтау министрлігінің «Денсаулық сақтауды ұйымдастыру және ақпараттандыру Орталық ғылыми-зерттеу институты» Федералды мемлекеттік бюджеттік мекемесі, e-mail: son@mednet.ru; ORCID:0000-0001-9309-2853.

Аталақова Жупар Ермековна – гуманитарлық білім магистрі, жарияланым белсенділігі бөлімінің бас маманы, «С. Д. Асфендияров атындағы Қазақ Ұлттық медицина университеті» КеАҚ; e-mail: e.zhupar@mail.ru; ORCID: 0000-0003-0025-7116.

ОБ АВТОРАХ

Юсупова Наргиза Сайдахметовна – магистр медицинских наук, менеджер по международному сотрудничеству, РГП на ПХВ «Казахский научный центр дерматологии и инфекционных заболеваний» Министерства здравоохранения Республики Казахстан; выпускник программы докторантурры по специальности «Общественное здравоохранение» НАО «Казахский Национальный Медицинский Университет имени С. Д. Асфендиярова»; e-mail: ns.yussupova@gmail.com; ORCID:0000-0003-1857-2800.

Турдалиева Ботагоз Сайтовна – доктор медицинских наук, профессор, заместитель директора по научной деятельности и стратегическому развитию, РГП на ПХВ «Казахский научный центр дерматологии и инфекционных заболеваний» Министерства здравоохранения Республики Казахстан; e-mail: botturd@gmail.com; ORCID:0000-0001-9884-0777.

Койков Виталий Викторович – доктор медицинских наук, проректор, НАО «Медицинский университет Астана»; e-mail: koykov@inbox.ru; ORCID:0000-0001-7668-5614.

Сон Ирина Михайловна – доктор медицинских наук, профессор, заместитель директора по науке и образованию, Федеральное государственное бюджетное учреждение «Центральный научно-исследовательский институт организаций и информатизации здравоохранения» Министерства здравоохранения Российской Федерации, e-mail: son@mednet.ru; ORCID:0000-0001-9309-2853.

Аталақова Жупар Ермековна – магистр гуманитарных знаний, главный специалист отдела публикационной активности, НАО «Казахский Национальный Медицинский Университет имени С. Д. Асфендиярова»; e-mail: e.zhupar@mail.ru; ORCID: 0000-0003-0025-7116.

ABOUT AUTHORS

Yussupova Nargiza Saidakhmetovna – master of Medical sciences, Manager of international cooperation, Kazakh scientific center of dermatology and infectious diseases of the Ministry of Health of the Republic of Kazakhstan; graduate of the doctoral program in the specialty «Public Health», NpJSC «Kazakh National Medical University named after S. D. Asfendiyarov»; e-mail: ns.yussupova@gmail.com; ORCID:0000-0003-1857-2800.

Turdaliyeva Botagoz Saitovna – doctor of Medical Sciences, Professor, Deputy Director of Science and Strategic Development, Kazakh scientific center of dermatology and infectious diseases of the Ministry of Health of the Republic of Kazakhstan; e-mail: botturd@gmail.com; ORCID:0000-0001-9884-0777.

Koikov Vitaliy Viktorovich – doctor of Medical Sciences, Vice-Rector, NpJSC «Astana Medical University»; e-mail: koykov@inbox.ru; ORCID:0000-0001-7668-5614.

Son Irina Mikhailovna – Doctor of Medical Sciences, Professor, Deputy Director of Science and Education, Federal Research Institute for Health Organization and Informatics of the Ministry of Health of the Russian Federation; e-mail: son@mednet.ru; ORCID:0000-0001-9309-2853

Atalykova Zhupar Yermekovna – master of Humanitarian Knowledge, Chief Specialist of the Department of Publication Activity, NpJSC «Kazakh National Medical University named after S. D. Asfendiyarov»; e-mail: e.zhupar@mail.ru; ORCID: 0000-0003-0025-7116.

Conflict of Interest. All authors declare that there is no potential conflict of interest requiring disclosure in this article.

Author Contributions. This article was prepared as part of the doctoral dissertation on the topic: "Scientifically Based Approaches to Improving Primary Health Care to Ensure the Operational Functions of Public Health" by the first and corresponding author, N.S. Yusupova. All other authors contributed equally to the development of the concept, implementation, data analysis, and writing of the article. We declare that this material has not been previously published and is not under consideration by any other publisher.

Funding. None.

Article submitted: 28.06.2025 year

Accepted for publication: 23.05. 2025 year